


## 5-13 Innovative Technique for Elucidating Diffusion through Materials

- Development of an Advanced Analysis Method for Quasielastic Neutron Scattering -




**Fig.5-34 Quasielastic neutron scattering of water**

Quasielastic scattering spectra obtained by a neutron scattering experiment. The signal is spread like a mound around the elastic scattering of zero energy transfer. Because more than one diffusion mode exists, the routes, speeds, and types of the modes are difficult to identify because their signals overlap. The red dot indicates the signal generated by the sample cell.

Diffusion is an important process in materials and living tissue. For instance, it is involved in the functional expression of proteins, and plays a role in the capabilities of ion and proton conductors, which are now being investigated as battery materials. Therefore, by elucidating diffusion mechanisms, we can control the functional expressions and capabilities of materials. Quasielastic neutron scattering (QENS) is among the few powerful techniques that can investigate the relationships between time and space in the diffusion motions of atoms and molecules. Therefore, QENS allows an in-depth study of diffusion. However, the correctness of diffusion models is difficult to demonstrate by conventional QENS analysis, as the model must be assumed prior to analysis, and therefore cannot naturally emerge from the results.

We have developed a versatile analysis method that requires no model assumptions. By this method, we can understand the model from the obtained results. We represent all diffusion processes of a material as a set of “simple diffusions.” In a “simple diffusion,” on average, the particles gradually spread out with a constant relaxation time. In this scenario, the diffusion process can be converted into the intensity



**Fig.5-35 Calculated mode distribution function of water**

Using the newly developed analysis, we computed the mode distribution function (MDF) of water. Each diffusion mode is independently represented. Consequently, we can directly evaluate the number of modes, and their types and speeds.

distribution of the relaxation time. We call this distribution the mode distribution function (MDF). The MDF directly provides the number of diffusion motions, and their types and speeds. The conversion to MDF adopts the maximum entropy method based on information theory.

We applied this new method to a diffusion study of water. QENS experiments were performed on AMATERAS at J-PARC. AMATERAS can measure over a wide area at high resolution and is ideally suited for this method. A QENS spectrum of liquid water is shown in Fig.5-34. The characteristics of the diffusion are not directly interpretable from this figure. However, the MDF obtained from the QENS spectrum (Fig.5-35) clearly shows three types of motions. In particular, an unsuspected intermediate motion is discovered. Therefore, the MDF reveals the detailed characters of diffusion motions. This result is important for understanding the overall picture of diffusion in water.

We hope that such diffusion studies will advance a wide range of research fields and uncover new knowledge that will benefit science and technology.

### Reference

Kikuchi, T. et al., Mode-Distribution Analysis of Quasielastic Neutron Scattering and Application to Liquid Water, *Physical Review E*, vol.87, no.6, 2013, p.062314-1-062314-8.